# Colonoscopy Preparation Instructions MAGNESIUM CITRATE (AM PROCEDURE)

## 1. THREE DAYS PRIOR TO YOUR PROCEDURE

- Please start a low residue/low fiber diet.
  - o Acceptable low residue/low fiber diet include:
    - Meat
    - Bacon
    - Fish
    - Chicken
    - Cheese
    - Eggs
    - Milk
    - Yogurt
    - Oatmeal

- Cream of
  - wheat
- Pancakes
- Waffles
- Cake
- Saltine
- Crackers
- White bread
- Pasta

- Rice
- Potato
- Squash
- Avocado
- Canned fruit
- Olive oil
- Butter
- Chocolate

# 2. THE ENTIRE DAY PRIOR TO YOUR PROCEDURE

- You may only have clear liquids
  - o Acceptable clear liquids include:
    - Beef or chicken broth
    - Jell-O or popsicles (NO RED, PURPLE OR ORANGE)
    - Sprite
    - 7-UP

- Ginger Ale
- Apple juice
- Water
- Tea or coffee (PLAIN, NO CREAM OR SUGAR)

- What you need: (2) Magnesium citrate bottles

  Dulcolax 5mg tablets (*four tablets*)
- Beginning at **7 PM** 
  - O Take the first 2 Dulcolax tablet
  - o Drink one bottle of the Magnesium citrate.

#### 3. THE DAY (MORNING) OF YOUR PROCEDURE

- Take the last 2 Dulcolax tablets **6 hours prior** to your arrival time.
- Take the last bottle of Magnesium citrate, 5 hours prior to your arrival time
  - \*\*Also, you may not have <u>ANYTHING</u> within 3 hours of your arrival time. \*\* Your procedure may be cancelled if you do so.
- 4. Arrange to take the day off of work since anesthesia will be administered. You MUST have a responsible adult driver present at the time of arrival and during your recovery. We cannot start the procedure without your driver present in the building. You will NOT be able to drive yourself home after the procedure. Uber, Lyft, Taxi, bus or other forms of "public transportation" are NOT acceptable means of returning home after the procedure. If you do not have an acceptable mode of transportation home after the procedure, your appointment will be cancelled and rescheduled to a different date.

**5.** All body piercings <u>MUST BE REMOVED</u> prior to arrival.

#### 6. MEDICATION MODIFICATIONS

### \*\*PLEASE DO NOT STOP TAKING YOUR ASPIRIN\*\*

- **Stop** Coumadin 5 days prior to the procedure
- Stop Pradaxa, Xarelto and Eliquis 2 days prior to the procedure
- Stop Plavix 7 days prior to the procedure
- **Stop** any anti-inflammatory medication 1 day prior to the procedure (including ibuprofen)
- Stop any arthritis medication 1 day prior to the procedure
- **DO NOT TAKE** insulin or any other oral diabetic medications the morning of the procedure. Be sure to check your blood sugar 4 hours prior to your appointment time. If your blood sugar is low, drink a glass of apple juice
- Stop Trulicity, Byetta, Saxenda, Victoza, Ozempic, Wegovy, Rybelsus, and Mounjaro ONE WEEK
   PRIOR to your procedure.
- Stop Canagliflorzin, Dapagliflozin, and Empagliflozin, THREE DAYS PRIOR to your procedure.
- Please take ½ dose Lantus the night before the procedure (if you are prescribed this)
- Continue to take any other prescription medications that are not listed above. They must be taken no later than 3 hours prior to your arrival time.

\*\*IF YOU ARE NOT SURE IF YOUR MEDICATIONS FALLS UNDER ANY OF THESE CATEGORIES, PLEASE CONTACT YOUR PHARMACY FOR FURTHER ASSISTANCE. \*\*

Your exam is scheduled on	at	am / pm
At	. Please arrive no later than	am / pm

# \*DO NOT ARRIVE EARILER THAN YOUR ARRIVAL TIME\*

YOUR PROCEDURE MAY BE CANCELLED AND RESCHEDULED IF YOU ARE LATE!

\*\*\*\*\*PLEASE BE FLEXIBLE. YOUR PROCEDURE TIME MAY VARY DEPENDING ON NUMEROUS FACTORS (cancellations, procedural delays, etc) \*\*\*\*

**Important**: It is important that you contact your insurance company <u>5 DAYS PRIOR</u> to your procedure date. You may have a co-payment the day of your procedure and are responsible to verify with your insurance in advance. Patients calling to cancel less than 2 business days due to a high co-payment may be subject to a \$150.00 cancellation fee.